

Drivers of medium- and large-bodied terrestrial mammal communities within a savannah–forest mosaic on Makushi and Wapichan customary lands in the Rupununi, Guyana

MATTHEW T. HALLETT^{*1,2,3} , FRANCESCO ROVERO^{3,4} 
NATHALIE VAN VLIET³ , FERNANDO M. LI^{5,6,7}, RHOMAYNE LI^{5,6}
JEREMY MELVILLE^{5,6}, ASAPH WILSON^{5,7}, HOWARD BARNABAS^{5,8}
ANDREW ALBERT^{5,9}, DEXTER DA SILVA⁵, ERIN EARL^{5,7}
OSWIN DAVID⁵ and SIMÓN QUINTERO^{3,10} 

Abstract The savannah–forest mosaic of the Rupununi region of Guyana is a dispersal corridor between large tracts of intact Guiana Shield forests and a subsistence hunting ground for Indigenous Makushi and Wapichan communities. We conducted a camera-trap survey at 199 sites across four major forested habitat types and used multi-species occupancy modelling to determine regional-scale drivers of mammalian occupancy at both species and community levels, accounting for imperfect detection. We detected 47 savannah- and forest-dwelling mammal species, with the occupancy of medium- and large-bodied terrestrial mammal species (community occupancy) positively related to per cent forest cover and negatively to the presence of gallery forest habitat. The occupancy of 15 of 30 species was positively related to forest cover, suggesting the importance of maintaining forested habitat within the broader mosaic comprising savannahs and intermediate habitats for sustaining maximum mammal diversity. Jaguar *Panthera onca* occupancy was associated with the presence of livestock, and giant anteater *Myrmecophaga tridactyla* occupancy was negatively associated with distance to the nearest road, both results of concern in relation to potential human–wildlife conflict. The probability of detecting terrestrial mammal species (community detectability) increased away from villages, as did the detectability of two large-bodied, hunted species, the lowland tapir *Tapirus*

terrestris and collared peccary *Pecari tajacu*, potentially indicating the negative effects of subsistence and commercial hunting in this savannah mosaic habitat. We use our findings to discuss how management strategies for hunting, fire, timber harvest and agriculture within Indigenous titled lands could help ensure the sustainability of these traditional livelihood activities.

Keywords Camera traps, community ecology, customary lands, Guyana, Indigenous science, multi-species Bayesian occupancy, Neotropical savannah, terrestrial mammals

The supplementary material for this article is available at doi.org/10.1017/S003060532400156X

Introduction

Amazonian savannahs are complex mosaics of open, seasonally flooded, natural grasslands surrounded by, and interspersed with, closed canopy forests with transition zones that support diverse vegetation communities, each of which provide food and cover and serve as pathways for the movement of unique mammal assemblages (Whitney & Davis, 1986; Risser, 1990; Ribeiro & Walter, 2008; Simon et al., 2009; Carmignotto et al., 2012; Maurin et al., 2014). Although habitat mosaics may support high overall diversity as a result of a mixture of vegetation communities, mosaics may also contain smaller or lower quality representations of distinct habitat types with which species are typically associated (Foster & Gaines, 1991). As a result of these species–area relationships, densities of large mammals may be naturally lower along edges or within forested fragments or corridors compared to their source habitats, although variation is driven by patch quality and connectivity within the landscape (Bowers & Matter, 1997).

Habitats within mosaics are defined by interactions between local climate, hydrology, herbivory and soil conditions (Hopkins, 1992; Staver et al., 2011), but fire seems to be the most important factor in maintaining Neotropical forest–savannah boundaries (Hoffmann et al.,

*Corresponding author, mhallett2320@ufl.edu

¹Department of Wildlife Ecology & Conservation, University of Florida, Gainesville, Florida, USA

²Conservation Department, Jacksonville Zoo & Gardens, Jacksonville, Florida, USA

³Center for International Forestry Research (CIFOR), Bogor, Indonesia

⁴Department of Biology, University of Florence, Sesto Fiorentino, Italy

⁵Sustainable Wildlife Management Programme (SWM) Guyana, Lethem, Guyana

⁶Rupununi Wildlife Research Unit, Lethem, Guyana

⁷South Rupununi Conservation Society, Wichabai Ranch, Guyana

⁸Caiman House Research Station, Yupukari Village, Guyana

⁹Keseé Ecolodge, Karasabai Village, Guyana

¹⁰Centro de Estudios Socioecológicos y Cambio Global, Instituto de Investigación de Recursos Biológicos Alexander von Humboldt, Bogotá, Colombia

Received 21 December 2023. Revision requested 2 April 2024.

Accepted 16 October 2024

2012). The presence of riparian habitat (i.e. gallery forest) may provide refuge from frequent wildfires (Prada & Marinho-Filho, 2004), seasonal access to resources (water, fruits) and forested dispersal corridors (Redford & Fonseca, 1986) for some species, and seasonal floods may also create adverse conditions and barriers to dispersal for others (Diniz-Filho et al., 2008). Although these generally open biomes comprise a large proportion of terrestrial landscapes (Furley, 2006), savannah biodiversity, and wildlife in particular, have been neglected in scientific research in comparison to tropical forests (Werneck, 2011).

Amazonian savannahs are threatened by agriculture and livestock (San José & Montes, 2001; Rosa et al., 2021), uncontrolled burning and infrastructure development (Mustin et al., 2017), all of which negatively impact mammal populations (de Marco et al., 2020; Romero-Muñoz et al., 2021). Reduction in both the quantity and quality of wildlife habitat also increases competition for resources, leading to conflict and retaliation for crop loss and depredation of livestock (de Souza et al., 2018). The advantageous combination of access to both savannah and forest resources also means that savannah–forest mosaics exist in close proximity to Indigenous communities that depend on wildlife for food and livelihoods (de Paula et al., 2017). Many Indigenous communities in the Amazon use fire to manage savannah grasses for livestock, to clear small patches of forest for traditional rotational agriculture, or for hunting (Mistry et al., 2016), but when poorly managed these practices may be a threat to ill-adapted tropical forest habitat. Hunting intensity, which has already reached unsustainable levels in many forested areas across the Neotropics (Redford, 1992; Fa & Peres, 2001; Wilkie et al., 2011), may also be higher in savannah–forest mosaics because of their proximity to human communities. Despite growing interest in the sustainability of hunting in the Neotropics, the response of mammal communities to hunting pressure within savannah and mosaic habitats remains largely understudied (Fernandes-Ferreira & Alves, 2017).

Here, we aimed to contribute to a better understanding of the ecological and anthropogenic drivers of the diversity and distribution of terrestrial medium- and large-bodied mammals (henceforth ‘large mammals’) across savannah–forest mosaic habitats. We conducted a region-wide camera-trap study in the hyper-seasonal Rupununi savannah of south-west Guyana. The area remains largely understudied despite comprising c. 5% of Amazon savannahs and supplying an important source of dietary protein for Indigenous communities. Our study was carried out within Makushi and Wapichan Indigenous community-owned (customary) lands where communities have maintained traditional management practices for millennia and more recently engaged in the development of management plans for the sustainable use of wildlife. We

hypothesized that the community of large mammals within this savannah–forest mosaic would be driven largely by forest cover and habitat type (with those habitats containing higher forest cover having higher species diversity and occupancy), and that anthropogenic factors would drive variation among species that are more susceptible to the effects of hunting, roads and livestock. Our assessment was carried out with the intention of informing management decisions in support of both traditional and modern local livelihoods in the Rupununi, while serving as a case study that may assist with the broader management of wildlife across Amazon savannahs.

Study area

The Rupununi savannahs are a low-lying, seasonally flooded, savannah–forest complex linked to Brazil’s Rio Branco savannahs (Montambault & Missa, 2002). They are divided into the North and South Rupununi by the Kanuku Mountains (Jansen-Jacobs & ter Steege, 2000), covering an area of c. 13,000 km². The Rupununi savannahs are contiguous with the Iwokrama International Centre for Rainforest Conservation & Development to the north, the Pakaraima Mountains and Gran Sabana to the west, and the Kanuku Mountains Protected Area to the east, which joins a vast expanse of intact Guiana Shield forest shared with Brazil, Suriname and French Guiana (Mittermeier et al., 1998).

The primary habitat types in the Rupununi, savannah and moist tropical forest, are principally determined by soil conditions, with savannah habitat occurring where trees cannot take root because a hard underlying clay layer limits penetration of their roots (Montambault & Missa, 2002). So-called bush islands are natural forest fragments that are scattered across the landscape on hilltops out of the reach of seasonal flood waters. Moist forest occurs on porous substrates along the slopes of hills and mountains, along rivers and in adjacent low-lying areas that receive nutrient runoff (Clark et al., 2001). The Rupununi savannahs are found at elevations of 120–150 m, the highest mountain peak is 1,067 m (Montambault & Missa, 2002), and gallery forests form along the edges of rivers and creeks. This region of Guyana experiences a single rainy season (May–August), and a longer dry season (September–April). Mean annual rainfall is 1,500–2,000 mm (Protected Areas Commission Guyana, 2015) and the main rivers rise by as much as 15 m during the peak of the rainy season, flooding low-lying forests and inundating adjacent savannahs.

The Rupununi region supports a low human population density (0.42 people/km²), with communities consisting of predominantly Indigenous Makushi and Wapichan people with mixed populations that include all of Guyana’s nine Indigenous groups, as well as Chinese-, Brazilian-, Afro-,

and Indo-Guyanese (Stone, 2002). The majority of Rupununi communities depend on subsistence activities for their livelihoods, with c. 20% of Rupununi households reporting that wild meat is their primary source of protein (along with 55% of households depending on fish; Luzar et al., 2012). The Rupununi region is accessed from Guyana's coastal capital via the 538 km long, largely unpaved, Georgetown–Lethem highway, and the remainder of the region is accessible via a network of unpaved roads constructed of a mix of laterite and sand (Paemelaere et al., 2023).

Methods

Data collection

Camera traps (Bushnell Trophy Cams 119837 and 119876; Bushnell, USA) were set 2–3 km apart, with a single camera at each trap location, set 30–40 cm from the ground in proximity to observed animal signs. Cameras were active continuously, with a 1-second delay between captures, recording the date and time with each 3-image sequence. Images of the same species at the same trap site within a period of 30 min were excluded to ensure that photo occasions were independent (O'Brien et al., 2003). No scents or lures were used, and all cameras were equipped with infrared flash.

Camera-trap sites were selected across an assortment of variables relevant at a range of scales: subregion, habitat type, trail type and distance to nearest water body, road and human settlement (Fig. 1). Key limiting factors for camera placement were community permission, accessibility and threat of flood and fire. Supplementary Material 1 provides a full description of criteria for site selection. Camera traps were set in clusters of 20 cameras, installed for 8–177 trap nights (mean 89.5), and then moved to a second location, with the same sampling effort. The initial survey period occurred during the rainy season, followed by a second during the dry season. Overall, 199 sites were surveyed from May 2019 to March 2020, totalling 16,382 trap nights, yielding 485,953 photographs over 24,800 occasions (Table 1).

Habitat type, per cent forest cover and distances to nearest trail, village and road were identified as relevant environmental and anthropogenic variables affecting the probability of occurrence of large mammals, with distance to nearest trail and village also used as potential covariates of detectability. Habitat type was determined at the site level using the definitions in Supplementary Material 1. Our predictions for the potential effect of these variables on occupancy and detectability are presented in Supplementary Material 2. Per cent forest cover was determined by placing a 1 km buffer around each camera trap and calculating the per cent forest cover from 30 m resolution global Landsat data (Hansen et al., 2013) within each buffer using the Tabulate

Area function. Distances to nearest village, road and trail were determined by calculating the Euclidean distance from camera-trap sites. All spatial analyses were conducted in *ArcMap 10.4.1* (Esri, 2016).

Prior to modelling, we verified that variables were not collinear using a Pearson's $r > 0.7$ threshold to indicate collinearity, and we did not detect any collinearity. Gallery forest and bush islands had a lower mean per cent forest cover than lowland and upland forest habitats (Supplementary Fig. 1).

Data analysis

We annotated camera-trap images manually and included in the analyses only mammals that are predominantly ground-dwelling and with a body mass of > 2 kg, which are the most likely to be detected by camera traps. Two pairs of species were similar in appearance and could not be differentiated reliably from photographs: *Dasypus guianensis* and *Dasypus kappleri*, as well as *Philander opossum* and *Metachirus nudicaudatus*; these were grouped and referred to as long-nosed armadillos and four-eyed opossums, respectively. An unknown species of cottontail rabbit that was detected during the survey is referred to as *Sylvilagus* sp.

We used a Bayesian hierarchical framework (Royle & Dorazio 2008) to model large mammal occupancy at the community (Ψ ; probability of occurrence for multiple species in each ecological unit) and species (ψ ; the probability of X species occurring in each ecological unit) level, with data augmentation to estimate species richness at the study area and camera-trap site level while accounting for the number of unobserved species (Dorazio et al., 2006). We fitted a single model with a limited number of covariates for which we hypothesized ecologically meaningful potential effects (Supplementary Table 2), following an established approach (e.g. Zipkin et al., 2010; Rich et al., 2016). We modelled intercepts and slopes of both regressions as species-specific random effects, drawn from a common prior distribution for the mammal community and with mean and variance hyper-parameters representing the average intercept and slopes across the observed mammal community and the variation among species. Details of the modelling framework are described in Supplementary Material 2.

We implemented the models in a Bayesian framework using *JAGS 4.3.0* (Plummer, 2003) in *R 4.1.2* (R Core Team, 2017) with the *R2jags 0.7-1* package (Su & Yajima, 2015). Our study design allowed us to account for imperfect detection and provided a robust estimate of detectability (p ; a measure of uncertainty in detection applied at the community or species level). We specified a Uniform (0, 1) prior distribution for the data augmentation parameter (Ω), a Normal (0, 0.01) prior distribution on




FIG. 1 Locations of camera traps across habitat types and the five subregions of the Rupununi Region of Guyana. (Readers of the printed journal are referred to the online article for a colour version of this figure.)

the logit scale for mean community detectability ($\mu_{\alpha 0}$) and occupancy ($\mu_{\beta 0}$) parameters, and for all the covariate coefficients on detectability and occupancy (α and β). We generated three parallel chains of 100,000 iterations with a burn-in of 20,000 iterations and thinning by 20 to derive summaries of parameter posterior distribution. Convergence of the Markov chains was satisfactory based on the Gelman–Rubin statistic, which was always ≤ 1.04 (Gelman & Rubin, 1992). We evaluated covariate effects by considering whether 90% Bayesian credible intervals (BCIs) encompassed zero. Community-level, species-specific and species trait-specific (body size and trophic guild) groupings were considered.

Results

Camera traps documented 47 mammal species in total across the survey area. Domestic species, people and 10

species of small mammals < 2 kg (Emmons, 1997) were excluded from subsequent analysis (Supplementary Table 1; Supplementary Plate 1).

Detections and estimated occupancy were heterogeneous across the 30 medium- and large-bodied native mammal species detected (Table 2). Eight species were disproportionally common, with an estimated occupancy > 0.5 and relative abundance index (RAI; calculated as the number of photo events divided by the number of trap nights, multiplied by 100) of 1.13–24.39: red-rumped agouti *Dasyprocta leporina*, ocelot *Leopardus pardalis*, common opossum *Didelphis marsupialis*, long-nosed armadillos, giant anteater *Myrmecophaga tridactyla*, red brocket deer *Mazama americana*, lowland paca *Cuniculus paca* and tayra *Eira barbara*. Six species were rarely detected, with an estimated occupancy < 0.1 and relative abundance index of 0.01–0.13: bush dog *Speothos venaticus*, capybara *Hydrochoerus hydrochaeris*, oncilla *Leopardus tigrinus*,

TABLE 1 Number of camera traps by habitat type, and number of trap nights, photographs and occasions, in the five subregions of the Rupununi Region of Guyana (Fig. 1).

	South Pakaraimas	North Rupununi	Central Rupununi	South Rupununi	Deep South Rupununi	Total
Camera traps: upland forest	26	0	22	0	0	48
Camera traps: lowland forest	0	18	5	28	4	55
Camera traps: bush island	14	12	2	2	20	40
Camera traps: gallery forest	2	10	11	8	15	46
Total camera traps	42	40	40	38	39	199
Number of trap nights	3,892	3,338	3,232	3,139	2,781	16,382
Number of photographs	64,961	66,379	76,072	102,432	176,109	485,953
Number of occasions	4,324	4,414	4,612	7,680	3,770	24,800

white-lipped peccary *Tayassu pecari*, southern naked-tailed armadillo *Cabassous unicinctus* and cottontail rabbit.

At the community level, the model estimated 30–34 species (90% BCI) to be present (mean $31 \pm \text{SD } 1.35$; Supplementary Table 2). The median estimated camera-trap site richness was 3–16 species. Plotting site richness against selected covariates shows that richness tended to increase with forest cover and distance from nearest village, and was highest in upland and lowland forest, lowest in gallery forests and intermediate in bush islands (Supplementary Fig. 1).

Mean community occupancy was best predicted by forest cover, with a significant positive effect, and by distance to nearest village, with a marginally significant positive effect. Distance to nearest road and livestock RAI had negative effects (Fig. 2; Supplementary Table 2). Community occupancy was broadly similar across habitat types, with the exception of gallery forest where it was significantly lower than average. The effects of forest cover, distance to nearest village, distance to nearest road and livestock relative abundance index on community occupancy are shown in Fig. 3. Mean community detectability was positively and significantly associated with distance to nearest village, and distance to nearest road had a negative effect (Fig. 2).

At the species level, forest cover was the most influential covariate, with 15 species whose occupancy was significantly and positively associated with it. Three species (white-tailed deer *Odocoileus cariacou*, giant anteater and crab-eating fox *Cerdocyon thous*) were significantly and negatively affected, and 12 species were not significantly affected (Supplementary Fig. 4). The negative effect of gallery forest habitat that was significant at community level was underlined by significant and negative effects at species level for red brocket deer, tayra, long-nosed armadillos and red-rumped agouti. The effect of this habitat on occupancy tended to be positive for the capybara and jaguarundi *Herpailurus yagouaroundi*. For covariates that were not significant at the community level, the species' effects varied greatly on a case-by-case basis. For example, the occupancy of five species (lowland tapir, four-eyed opossums, collared peccary, red acouchi *Myoprocta acouchy* and Amazonian brown brocket deer *Mazama nemorivaga*) were significantly and positively associated with distance to nearest village. Similarly, puma *Puma concolor* occupancy was positively associated, and crab-eating raccoon *Procyon cancrivorus*, giant anteater, common opossum and red-rumped agouti were negatively associated with distance to nearest road. Jaguar occupancy was positively and significantly associated, and lowland paca and long-nosed armadillo negatively associated with livestock RAI (Fig. 4).

When we explored the magnitude of the effect of covariates on species traits, we found that forest cover,

TABLE 2 Naïve occupancy (Ψ), number of photographic occasions, relative abundance index (RAI; calculated as the number of photo events divided by the number of trap nights, multiplied by 100), occupancy ($\Psi \pm \text{SD}$) and variables with statistically significant effects, with direction of relationship (+ or -) for each of the 30 medium- and large-bodied mammal species detected, ordered by decreasing Ψ .

Species	Naïve Ψ	Occasions	RAI	$\Psi \pm \text{SD}$	Significant variables ¹
Red-rumped agouti <i>Dasyprocta leporina</i>	0.78	4,258	24.39	0.78 $\pm$ 0.03	FC+, DR-, GF-
Ocelot <i>Leopardus pardalis</i>	0.56	294	1.68	0.70 $\pm$ 0.04	FC+
Common opossum <i>Didelphis marsupialis</i>	0.63	590	3.37	0.67 $\pm$ 0.03	FC+, DR-
Long-nosed armadillos <i>Dasypus novemcinctus</i> / <i>Dasypus kappleri</i>	0.60	821	4.70	0.62 $\pm$ 0.03	FC+, LV-, GF-
Giant anteater <i>Myrmecophaga tridactyla</i>	0.49	328	1.87	0.56 $\pm$ 0.04	FC-, DR-
Red brocket deer <i>Mazama americana</i>	0.52	602	3.44	0.55 $\pm$ 0.03	FC+, GF-
Lowland paca <i>Cuniculus paca</i>	0.54	1,008	5.77	0.55 $\pm$ 0.03	FC+, LV-
Tayra <i>Eira barbara</i>	0.41	198	1.13	0.52 $\pm$ 0.04	FC+, GF-
Margay <i>Leopardus wiedii</i>	0.24	72	0.41	0.46 $\pm$ 0.08	FC+
Greater grison <i>Galictis vittata</i>	0.05	9	0.05	0.38 $\pm$ 0.21	
Puma <i>Puma concolor</i>	0.23	79	0.45	0.36 $\pm$ 0.05	FC+, DR+
Jaguarundi <i>Herpailurus yagouaroundi</i>	0.17	47	0.27	0.35 $\pm$ 0.07	
South American coati <i>Nasua nasua</i>	0.23	102	0.58	0.32 $\pm$ 0.05	
Collared peccary <i>Pecari tajacu</i>	0.28	182	1.04	0.32 $\pm$ 0.03	DV+, FC+
White-tailed deer <i>Odocoileus cariacou</i>	0.25	126	0.72	0.31 $\pm$ 0.04	FC-
Four-eyed opossums <i>Philander opossum</i> / <i>Metachirus nudicaudatus</i>	0.25	262	1.50	0.26 $\pm$ 0.03	DV+, FC+, DR-
Jaguar <i>Panthera onca</i>	0.13	38	0.22	0.24 $\pm$ 0.06	LV+
Crab-eating raccoon <i>Procyon cancrivorus</i>	0.16	65	0.37	0.24 $\pm$ 0.04	DR-
Red acouchi <i>Myoprocta acouchy</i>	0.23	1,277	7.31	0.23 $\pm$ 0.03	DV+, FC+
Lowland tapir <i>Tapirus terrestris</i>	0.18	132	0.76	0.21 $\pm$ 0.03	DV+, FC+
Southern tamandua <i>Tamandua tetradactyla</i>	0.13	47	0.27	0.20 $\pm$ 0.04	
Giant armadillo <i>Priodontes maximus</i>	0.08	24	0.14	0.19 $\pm$ 0.07	FC+
Amazonian brown brocket deer <i>Mazama nemorivaga</i>	0.16	156	0.89	0.18 $\pm$ 0.02	FC+
Crab-eating fox <i>Cerdocyon thous</i>	0.10	63	0.36	0.11 $\pm$ 0.02	FC-
Bush dog <i>Speothos venaticus</i>	0.01	2	0.01	0.09 $\pm$ 0.10	
Capybara <i>Hydrochoerus hydrochaeris</i>	0.06	23	0.13	0.08 $\pm$ 0.02	
Oncilla <i>Leopardus tigrinus</i>	0.03	8	0.05	0.08 $\pm$ 0.05	
White-lipped peccary <i>Tayassu pecari</i>	0.02	4	0.02	0.06 $\pm$ 0.06	
Southern naked-tailed armadillo <i>Cabassous unicinctus</i>	0.02	13	0.07	0.03 $\pm$ 0.02	
Cottontail rabbit <i>Sylvilagus</i> sp.	0.02	18	0.10	0.02 $\pm$ 0.01	

¹DV, distance to nearest village; FC, forest cover; LV, livestock RAI; DR, distance to nearest road; GF, presence of gallery forest.




FIG. 2 Effect size plot for the influence of covariates on (a) community occupancy and (b) community detectability. The effects of habitat types are relative to the baseline level (bush islands). Segments show 90% (thick) and 50% (thin) Bayesian credible intervals, with black segments indicating those for which 90% BCI does not overlap zero.

distance to nearest village and livestock RAI showed no pattern in relation to body mass, and distance to nearest road had a positive effect only on the largest species (Supplementary Fig. 2). Forest cover appeared least influential for the insectivores compared to other guilds; herbivores/frugivores and omnivores were more influenced by distance to nearest village than carnivores and insectivores; livestock RAI influenced the carnivores and omnivores more than the other guilds, and there was no clear pattern for distance to nearest road (Supplementary Fig. 3).

Discussion

Our study confirms the presence of 30 native, terrestrial medium- and large-bodied mammal species in the Rupununi, a diversity comparable to that of other rich neotropical savannah–forest systems, such as the Pantanal (Trolle, 2003). Average mammal community occupancy was similar across habitat types, except for within gallery forest habitat where it was significantly lower, and showed a significant positive association with forest cover, a marginally significant positive association with distance to nearest village, and negative associations with distance to nearest road and livestock detection rate. As hypothesized, species-specific responses were variable and primarily linked to human activities.

Forest cover

Species richness increased with per cent forest cover, which was also the most important driver of community occupancy overall, and had a significant positive

relationship with 15 of the species. The positive association is expected for semi-arboreal species such as the margay, tayra, common opossum and four-eyed opossum, which depend on tree canopies to hunt, forage or rest. A wide variety of herbivores/frugivores (tapir, brocket deer, collared peccary, paca, agouti, acouchi) that depend on the leaves, fruits and seeds of forest trees, palms and vines also had a significant positive relationship with forest cover. Puma occupancy was also positively correlated with per cent forest cover; this field primarily preys on mammalian herbivore/frugivore species (Foster et al., 2010) that are associated with forested areas.

The occupancy of typically savannah-dwelling species (the crab-eating fox, white-tailed deer and giant anteater) had significant negative relationships with per cent forest cover. Although the former two are more limited in their distribution to open savannahs and forest edges, the latter readily inhabits intact forests but maintains higher population densities in open habitats with lower predator abundance (Quiroga et al., 2016). Another six species (the southern naked-tailed armadillo, southern tamandua, capybara, crab-eating raccoon, South American coati and jaguarundi) had weak negative relationships with forest cover, showing their ability to utilize a variety of closed and open canopy habitats. All of these species feed on insects (forest cover appeared least influential for insectivores relative to other guilds), grasses or a generalized omnivorous diet.

Habitat type

Large mammal community occupancy was similar across habitat types, with the exception of a significant negative




FIG. 3 Bivariate effects on community occupancy of (a) distance to nearest village, (b) per cent forest cover, (c) livestock relative abundance index (RAI), (d) distance to nearest road and (e) habitat, as estimated from camera trapping at 195 sites in Guyana (Fig. 1).

correlation with gallery forest habitat. Although individuals of some species may make seasonal use of gallery forests for dispersal or accessing water or palm fruits, gallery forest provides the least forest cover and seasonal flooding may limit the dispersal of some species, likely driving the negative relationship with most species (28 of 30). Thus, species with a clear preference for terra firme forest, such as the red brocket deer, tayra, red-rumped agouti and long-nosed armadillo, had a significant negative relationship with gallery forest. The capybara and jaguarundi were the only species with positive relationships with gallery forest habitat. The capybara eats riverside grasses and is semi-aquatic, characteristics suitable for this seasonally inundated environment (Moreira et al., 2012). The jaguarundi is a good swimmer (Escobar-Anleu et al., 2020) and uses open habitats (Giordano, 2016) to avoid competition with larger felids (Caso, 2013).

Bush islands had the highest mean community occupancy (Fig. 3e), potentially because of their accessibility to both savannah and forest-dwelling species. Although savannah species may make use of the cover provided by bush islands, the size of each individual bush island and its proximity to other bush islands within a network of forested stepping stones spread out across the landscape is likely the most important factor in predicting their use by forest-dwelling species. Increased farming, hunting and

harvesting of large trees to supply growing village populations has resulted in observed declines in the size of, and diversity supported by, many bush islands. The high community occupancy of bush islands makes the management of these habitats, especially larger islands and those in close proximity to intact forest, particularly important.

Upland and lowland forest habitats had the highest mean forest cover, the least variation in forest cover, and the highest species richness among the sites sampled. These habitats are largely intact, managed for generations by the traditional management systems of the Indigenous communities of the Rupununi region. With their proximity to open savannah, lowland forests may vary in structure and composition and this forested ecotone also experiences the region's highest rate of degradation and loss, primarily as a result of uncontrolled burning (Global Forest Watch, 2022). This issue is of growing concern because of more recent increases in both the frequency and duration of savannah fires, the result of the combination of a hotter, drier climate and loss of traditional fire management practices (Mistry et al., 2016).

Distance to nearest village

Species richness tended to increase, and mammal community occupancy showed a marginally significant, positive

correlation, with distance to nearest village. We believe these trends are likely driven by the effect of subsistence and commercial hunting, which provides a primary source of protein and income for many households and predominantly occurs 6–12 km from village centres (Read et al., 2010). We cannot completely discount the effect of environmental noise associated with human settlements in reducing abundance (Shannon et al., 2016), as mean community detectability was also positively and significantly associated with distance to the nearest village. Studies of hunted areas within intact forest habitats experiencing lower hunting intensities (survey locations > 20 km from any village) have detected shifts in activity patterns away from peak times of human activity, but only detected weak effects on occupancy (Roopsind et al., 2017; Hallett et al., 2019). The community-level effects detected in our study are likely the result of a combination of naturally reduced densities of species in savannah–forest mosaics (when compared to intact forest), along with proximity to villages, which adds the combined effects of hunting, fire, farming, roads and livestock.

At the species level, the occupancy of two highly sought-after game species, the lowland tapir and collared peccary, had significant positive relationships with distance to nearest village. A number of forest-dwelling game species (the red and brown brocket deer, white-lipped peccary, lowland paca, and giant and long-nosed armadillos) also had weak positive relationships with distance to nearest village. Puma occupancy also had a marginally significant positive relationship with distance to nearest village, indicating that the effect of hunting detected in several important prey species is also having cascading effects on food chains. Pumas are known to feed on larger prey than other sympatric carnivores (Foster et al., 2010) and their abundance decreases with prey abundance (Laundré et al., 2007). Although the species-level effects may suggest that hunting is currently only negatively affecting a few large-bodied species preferred by hunters, the positive correlation between distance to nearest village and the occupancy of a suite of smaller-bodied or less preferred game species and an adaptable species like the puma is of concern with respect to the sustainability of current levels of subsistence and commercial hunting within the Rupununi savannah–forest mosaic.

Distance to nearest road

Giant anteater, red-rumped agouti, crab-eating raccoon and common and four-eyed opossums had significant negative relationships with distance to nearest road, likely driven by opportunities to forage on young plants and insects, scavenge on roadkill and discarded food, and drink from roadside ditches. However, a number of large-bodied

game species (the giant armadillo, white-lipped peccary, lowland tapir and brown brocket deer) and the puma (primary predator of large mammals) all had positive relationships (significant for the puma) with distance to roads, indicating the negative impact of road access on species that may be hunted more efficiently from roads (Ramos-Robles et al., 2013) and/or are susceptible to being struck by vehicles (Cáceres et al., 2010). Giant anteaters had a significant negative relationship with distance to roads, which raises concern because they are particularly vulnerable to collisions with vehicles (Paemelaere et al., 2023) and the expansion of road networks decreases the suitability of their habitat (Pinto et al., 2018). Proactive mitigation measures in the design of the current and proposed upgrades to the Georgetown–Lethem highway would help to reduce the impact of roads on this threatened species.

Jaguars and livestock

The occupancy of carnivores overall had a positive relationship with livestock RAI, and jaguar occupancy did not have a significant correlation with any other variable. Despite their popular association with tropical forests and ungulate prey, jaguars are highly adaptable opportunistic predators that inhabit a variety of habitats from moist forests to swamps, dry forests, scrub and grasslands, and prey on a wide range of mammals, reptiles and birds (Quigley et al., 2017). Jaguars are regularly implicated in depredation of livestock (de Azevedo & Murray, 2007; Foster et al., 2010) and depredation rates tend to increase with reduction in natural prey (Burgas et al., 2014). Prey densities in our study area were noticeably lower than in the neighbouring Kanuku Mountains (Hallett et al., 2019) or Iwokrama forest (Harris et al., 2023), probably reduced by the combination of natural and anthropogenic factors discussed above. Rupununi livestock are mainly managed in herds that range free over large areas, a key factor increasing depredation and conflict (de Azevedo & Murray, 2007; Montalvo et al., 2016). Although depredation rates in the Rupununi savannahs are not as high as elsewhere in the Neotropics, responses tend to be harsh, and retaliatory killing is the primary threat to jaguar populations (Hallett et al., 2024). This association raises a concern that the Rupununi savannah–forest mosaic may be an ecological trap for jaguar populations, and warrants further research.

Conclusions

The Rupununi Region of Guyana has a rich and dynamic Amazonian savannah, but also faces a diverse collection of threats that are challenging to manage. Our study provides

baseline data from a collaboration designed and implemented between Indigenous scientists and external partners. Indigenous communities in Guyana maintain ownership of their lands, and thus data-driven management strategies, such as the protection of sacred sites, improved control of hunters from outside the region, and the promotion of traditional hunting methods, could improve sustainability within these vital hunting areas. Management strategies to reduce or mitigate fire, timber harvest and agriculture within bush islands and lowland forest edges would also benefit important forested habitats within the mosaic. Implementation of improved livestock management and roadkill mitigation structures would help reduce conflict with species of conservation concern, the jaguar and giant anteater, respectively.

Author contributions Study design, data collection: MH, NV, FL, RL, JM, AW, HB, AA, DS, EE, OD; data analysis: FR, MH, SQ; writing: MH, NV, FR, SQ.

Acknowledgements We are grateful to the leadership and residents of Sustainable Wildlife Management Programme partner communities for their kindness, hospitality, generosity, cooperation, support and participation, without which this study could not have been carried out. This activity is part of the Sustainable Wildlife Management Programme, an Initiative of the Organization of African, Caribbean and Pacific States, which is funded by the European Union implemented through a consortium partnership that includes the Centre for International Forestry Research, the Food and Agriculture Organization of the United Nations, the Wildlife Conservation Society and the French Agricultural Research Centre for International Development.

Conflicts of interest None.

Ethical standards This research abided by the *Oryx* guidelines on ethical standards. Research protocols were reviewed, exempted from full review because the study was observational in nature and posed minimal risk, and approved by the Institutional Animal Care and Use Committee (IACUC) at the University of Florida. A free, prior and informed consent process facilitated by the Sustainable Wildlife Management Guyana Programme resulted in letters of consent signed by the 27 Indigenous communities that participated in this study. Research permits were approved by the Ministry of Amerindian Affairs and Environmental Protection Agency (#250219 BR01) in Guyana. Any photographs of people inadvertently obtained by camera traps were noted (i.e. recorded as people being present, without any further details) and then immediately and permanently deleted from the dataset.

Data availability Participating Indigenous communities expressed their preference to restrict access to the data, to protect community resources and intellectual property.

References

- BOWERS, M.A. & MATTER, S.F. (1997) Landscape ecology of mammals: relationships between density and patch size. *Journal of Mammalogy*, **78**, 999–1013.
- BURBAS, A., AMIT, R. & LOPEZ, B.C. (2014) Do attacks by jaguars *Panthera onca* and pumas *Puma concolor* (Carnivora: Felidae) on livestock correlate with species richness and relative abundance of wild prey? *Revista de Biología Tropical*, **62**, 1459–1467.
- CÁCERES, N.C., HANNIBAL, W., FREITAS, D.R., SILVA, E.L., ROMAN, C. & CASELLA, J. (2010) Mammal occurrence and roadkill in two adjacent ecoregions (Atlantic Forest and Cerrado) in south-western Brazil. *Zoologia (Curitiba)*, **27**, 709–717.
- CARMIGNOTTO, A.P., VIVO, M.D. & LANGGUTH, A. (2012) Mammals of the Cerrado and Caatinga: distribution patterns of the tropical open biomes of Central South America. In *Bones, Clones and Biomes: The History and Geography of Recent Neotropical Mammals* (eds B.D. Patterson & L.P. Costa), pp. 307–350. University of Chicago Press, Chicago, Illinois, USA.
- CASO, A. (2013) *Spatial differences and local avoidance of ocelot (Leopardus pardalis) and jaguarundi (Puma yagouaroundi) in northeast Mexico*. Doctoral dissertation, Texas A&M University-Kingsville, Texas, USA.
- CLARK, D.A., BROWN, S., KICKLIGHTER, D.W., CHAMBERS, J.Q., THOMLINSON, J.R., NI, J. & HOLLAND, E.A. (2001) Net primary production in tropical forests: an evaluation and synthesis of existing field data. *Ecological Applications*, **11**, 371–384.
- DE AZEVEDO, F.C.C. & MURRAY, D.L. (2007) Evaluation of potential factors predisposing livestock to predation by jaguars. *Journal of Wildlife Management*, **71**, 2379–2386.
- DE MARCO, P., VILLÉN, S., MENDES, P., NÓBREGA, C., CORTES, L., CASTRO, T. & SOUZA, R. (2020) Vulnerability of Cerrado threatened mammals: an integrative landscape and climate modeling approach. *Biodiversity and Conservation*, **29**, 1637–1658.
- DE PAULA, M., XERENTE, V. & PEZZUTI, J. (2017) Hunting and monitoring: community-based research in Xerente Indigenous Land, Brazilian Cerrado. *Human Ecology Review*, **23**, 23–44.
- DE SOUZA, J.C., DA SILVA, R.M., GONÇALVES, M.P.R., JARDIM, R.J.D. & MARKWITH, S.H. (2018) Habitat use, ranching, and human-wildlife conflict within a fragmented landscape in the Pantanal, Brazil. *Biological Conservation*, **217**, 349–357.
- DINIZ-FILHO, J.A.F., BINI, L.M., VIEIRA, C.M., BLAMIRE, D., TERRIBLE, L.C., BASTOS, R.P. et al. (2008) Spatial patterns of terrestrial vertebrate species richness in the Brazilian Cerrado. *Zoological Studies*, **47**, 146.
- DORAZIO, R.M., ROYLE, J.A., SODERSTROM, B. & GLIMSKAR, A. (2006) Estimating species richness and accumulation by modeling species occurrence and detectability. *Ecology*, **87**, 1650–1658.
- EMMONS, L.H. (1997) *Neotropical Rainforest Mammals* (2nd ed.). The University of Chicago Press, Chicago, Illinois, USA and London, UK.
- ESRI (2016) *ArcGIS Release 10.4.1*. Environmental Systems Research Institute Redlands, California, USA.
- ESCOBAR-ANLEU, B.I., QUIRÓNEZ-GUZMÁN, J.M. & HERNÁNDEZ-GÓMEZ, S. (2020) First swim record of *Herpailurus yagouaroundi* in Guatemala. *Therya Notes*, **1**, 29–33.
- FA, J.E. & PERES, C.A. (2001) Game vertebrate extraction in African and Neotropical forests: an intercontinental comparison. In *Conservation of Exploited Species* (eds J. Reynolds, G. Mace, J. G. Robinso & K. Redford), pp. 203–241. Cambridge University Press, Cambridge, UK.
- FERNANDES-FERREIRA, H. & ALVES, R.R.N. (2017) The researches on the hunting in Brazil: a brief overview. *Ethnobiology and Conservation*, **6**, 6.
- FOSTER, J. & GAINES, M.S. (1991) The effects of a successional habitat mosaic on a small mammal community. *Ecology*, **72**, 1358–1373.
- FOSTER, R.J., HARMSSEN, B.J. & DONCASTER, C.P. (2010) Habitat use by sympatric jaguars and pumas across a gradient of human disturbance in Belize. *Biotropica*, **42**, 724–731.
- FURLEY, P. (2006) Tropical savannas. *Progress in Physical Geography*, **30**, 105–121.

- GELMAN, A. & RUBIN, D.B. (1992) A single series from the Gibbs sampler provides a false sense of security. *Bayesian Statistics*, 4, 625–631.
- GIORDANO, A.J. (2016) Ecology and status of the jaguarundi *Puma yagouaroundi*: A synthesis of existing knowledge. *Mammal Review*, 46, 30–43.
- Global Forest Watch. (2022) Guyana. globalforestwatch.org/dashboards/country/GUY [accessed 3 February 2023].
- HALLETT, M., KAWALL, S., PIERRE, M., MELVILLE, A., KENYON, M., DE FREITAS, K. et al. (2024) Coexistence between people, livestock, and carnivores in the Rupununi. In *Wildlife and People in the Rupununi: A Comprehensive Assessment Based on Science and Local Knowledge* (eds N. van Vliet, N. Ponta, S. LeBel, L. McKenna & O. David), pp. 135–152. FAO, CIRAD, CIFOR-ICRAF, and WCS, Rome, Italy. [dx.doi.org/10.4060/cdo660en](https://doi.org/10.4060/cdo660en).
- HALLETT, M.T., KINAHAN, A.A., MCGREGOR, R., BAGGALLAY, T., BABB, T., BARNABUS, H. et al. (2019) Impact of low-intensity hunting on game species in and around the Kanuku Mountains Protected Area, Guyana. *Frontiers in Ecology and Evolution*, 7, 412.
- HANSEN, M.C., POTAPOV, P.V., MOORE, R., HANCHER, M., TURUBANOVA, S.A.A., TYUKAVINA, A. et al. (2013) High-resolution global maps of 21st-century forest cover change. *Science*, 342, 850–853.
- HARRIS, A.E., HALLETT, M.T., DAVIS, M., CARTER, M., SINGH, D., ROOPSIND, A. et al. (2023) Use of logging roads by terrestrial mammals in a responsibly managed neotropical rainforest in Guyana. *Forest Ecology and Management*, 548, 121401.
- HOFFMANN, W.A., GEIGER, E.L., GOTSCH, S.G., ROSSATTO, D.R., SILVA, L.C., LAU, O.L. et al. (2012) Ecological thresholds at the savanna-forest boundary: how plant traits, resources and fire govern the distribution of tropical biomes. *Ecology Letters*, 15, 759–768.
- HOPKINS, B. (1992) Ecological processes at the forest-savanna boundary. In *Nature and Dynamics of Forest-Savanna Boundaries* (eds P.A. Furley, J. Proctor & J.A. Ratter), pp. 21–33. Chapman & Hall, London, UK.
- JANSEN-JACOBS, M.J. & TER STEEGE, H. (2000) Southwest Guyana: A complex Mosaic of Savannahs and Forests. In *Plant diversity in Guyana: with recommendations for a National Protected Area Strategy* (ed H. ter Steege), pp. 147–157. Tropenbos International, the Netherlands.
- LAUNDRÉ, J.W., HERNÁNDEZ, L. & CLARK, S.G. (2007) Numerical and demographic responses of pumas to changes in prey abundance: testing current predictions. *The Journal of Wildlife Management*, 71, 345–355.
- LUZAR, J.B., SILVIUS, K.M. & FRAGOSO, J.M. (2012) Church affiliation and meat taboos in indigenous communities of Guyanese Amazonia. *Human Ecology*, 40, 833–845.
- MAURIN, O., DAVIES, T.J., BURROWS, J.E., DARU, B.H., YESSOUFOU, K., MUASYA, A.M. et al. (2014) Savanna fire and the origins of the ‘underground forests’ of Africa. *New Phytologist*, 204, 201–214.
- MISTRY, J., BILBAO, B.A. & BERARDI, A. (2016) Community owned solutions for fire management in tropical ecosystems: case studies from Indigenous communities of South America. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 371, 20150174.
- MITTERMEIER, R.A., MYERS, N., THOMSEN, J.B., DA FONSECA, G.A. & OLIVIERI, S. (1998) Biodiversity hotspots and major tropical wilderness areas: approaches to setting conservation priorities. *Conservation Biology*, 12, 516–520.
- MONTALVO, V., ALFARO, L., SAENZ, C., CRUZ, J., FULLER, T.K. & CARRILLO, E. (2016) Factors affecting jaguar and puma predation on livestock in Costa Rica. *Wildlife Biology in Practice*, 12, 32–42.
- MONTAMBAULT, J.R. & MISSA, O. (2002) *A Biodiversity Assessment of the Eastern Kanuku Mountains, Lower Kwitaro River, Guyana*. Rapid Assessment Program Bulletin of Biological Assessment 26. Conservation International, Washington, DC, USA. pdf.usaid.gov/pdf_docs/Pnadb868.pdf [accessed 20 January 2023].
- MOREIRA, J.R., FERRAZ, K.M.P., HERRERA, E.A. & MACDONALD, D.W. (2012) *Capybara: Biology, use and Conservation of an Exceptional Neotropical Species*. Springer, Berlin, Germany.
- MUSTIN, K., CARVALHO, W.D., HILÁRIO, R.R., COSTA-NETO, S.V., SILVA, C., VASCONCELOS, I.M. et al. (2017) Biodiversity, threats and conservation challenges in the Cerrado of Amapá, an Amazonian savanna. *Nature Conservation*, 22, 107–127.
- O'BRIEN, T.G., KINNAIRD, M.F. & WIBISONO, H.T. (2003) Crouching tigers, hidden prey: Sumatran tiger and prey populations in a tropical forest landscape. *Animal Conservation*, 6, 131–139.
- PAEMELAERE, E.A., MEJÍA, A., QUINTERO, S., HALLETT, M., LI, F., WILSON, A. et al. (2023) The road towards wildlife friendlier infrastructure: mitigation planning through landscape-level priority settings and species connectivity frameworks. *Environmental Impact Assessment Review*, 99, 107010.
- PINTO, F.A., BAGER, A., CLEVINGER, A.P. & GRILO, C. (2018) Giant anteater (*Myrmecophaga tridactyla*) conservation in Brazil: analysing the relative effects of fragmentation and mortality due to roads. *Biological Conservation*, 228, 148–157.
- PLUMMER, M. (2003) *JAGS: A Program for Analysis of Bayesian Graphical Models Using Gibbs Sampling*. Proceedings of the 3rd International Workshop on Distributed Statistical Computing, Vienna, Austria.
- PRADA, M. & MARINHO-FILHO, J. (2004) Effects of fire on the abundance of Xenarthrans in Mato Grosso, Brazil. *Austral Ecology*, 29, 568–573.
- PROTECTED AREAS COMMISSION GUYANA (2015) Kanuku Mountains Protected Area: Management Plan 2015–2019. Protected Areas Commission, Georgetown, Guyana. pac.gov.gy/wp-content/uploads/2022/10/KMPA-Plan.pdf [accessed December 2024].
- QUIGLEY, H., FOSTER, R., PETRACCA, L., PAYAN, E., SALOM, R. & HARMSSEN, B. (2017) *Panthera onca*. In *The IUCN Red List of Threatened Species* 2017. [dx.doi.org/10.2305/IUCN.UK.2017-3.RLTS.T15953A50658693.en](https://doi.org/10.2305/IUCN.UK.2017-3.RLTS.T15953A50658693.en).
- QUIROGA, V.A., NOSS, A.J., BOAGLIO, G.I. & DI BITETTI, M.S. (2016) Local and continental determinants of giant anteater (*Myrmecophaga tridactyla*) abundance: biome, human and jaguar roles in population regulation. *Mammalian Biology*, 81, 274–280.
- R CORE TEAM (2017) *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- RAMOS-ROBLES, M., GALLINA, S. & MANDUJANO, S. (2013) Habitat and human factors associated with white-tailed deer density in the tropical dry forest of Tehuacan-Cuicatlan Biosphere Reserve, Mexico. *Tropical Conservation Science*, 6, 70–86.
- READ, J.M., FRAGOSO, J.M., SILVIUS, K.M., LUZAR, J., OVERMAN, H., CUMMINGS, A. et al. (2010) Space, place, and hunting patterns among indigenous peoples of the Guyanese Rupununi region. *Journal of Latin American Geography*, 213–243.
- REDFORD, K.H. (1992) The empty forest. *BioScience*, 42, 412–422.
- REDFORD, K.H. & FONSECA, G.A.B. (1986) The role of gallery forests in the zoogeography of the Cerrado's non-volant Mammalian Fauna. *Biotropica*, 18, 126–135.
- RIBEIRO, J.F. & WALTER, B.M.T. (2008) As principais fitofisionomias do bioma Cerrado. *Cerrado: Ecologia e Flora*, 1, 151–212.
- RICH, L.N., MILLER, D.A., ROBINSON, H.S., MCNUTT, J.W. & KELLY, M.J. (2016) Using camera trapping and hierarchical occupancy modelling to evaluate the spatial ecology of an African mammal community. *Journal of Applied Ecology*, 53, 1225–1235.

- RISSE, P.G. (1990) The ecological importance of land-water ecotones. *The Ecology and Management of Aquatic-Terrestrial Ecotones*, 4, 7–21.
- ROMERO-MUÑOZ, A., FANDOS, G., BENÍTEZ-LÓPEZ, A. & KUEMMERLE, T. (2021) Habitat destruction and overexploitation drive widespread declines in all facets of mammalian diversity in the Gran Chaco. *Global Change Biology*, 27, 755–767.
- ROOPSIND, A., CAUGHLIN, T.T., SAMBHU, H., FRAGOSO, J. & PUTZ, F.E. (2017) Logging and indigenous hunting impacts on persistence of large Neotropical animals. *Biotropica*, 49, 565–575.
- ROSA, T.F., CAMAROTA, F., ZUANON, L.A., TITO, R., MARAVALHAS, J.B., POWELL, S. & VASCONCELOS, H.L. (2021) The effects of high-severity fires on the arboreal ant community of a Neotropical savanna. *Oecologia*, 196, 951–961.
- ROYLE, J.A. & DORAZIO, R.M. (2008). *Hierarchical modeling and inference in ecology: the analysis of data from populations, metapopulations and communities*. Elsevier, Cambridge, Massachusetts, USA.
- SAN JOSÉ, J.J., MONTES, R.A. (2001) Management effects on carbon stocks and fluxes across the Orinoco savannas. *Forest Ecology and Management*, 150, 293–311.
- SHANNON, G., MCKENNA, M.F., ANGELONI, L.M., CROOKS, K.R., FRISTRUP, K.M., BROWN, E. et al. (2016) A synthesis of two decades of research documenting the effects of noise on wildlife. *Biological Reviews*, 91, 982–1005.
- SIMON, M.F., GREYER, R., DE QUEIROZ, L.P., SKEMA, C., PENNINGTON, R.T. & HUGHES, C.E. (2009) Recent assembly of the Cerrado, a neotropical plant diversity hotspot, by in situ evolution of adaptations to fire. *Proceedings of the National Academy of Sciences*, 106, 20359–20364.
- STAYER, A.C., ARCHIBALD, S. & LEVIN, S.A. (2011) The global extent and determinants of savanna and forest as alternative biome states. *Science*, 334, 230–232.
- STONE, S. (2002) *Kanuku Mountains Protected Area Process: Community Resource Evaluation Master Report*. Conservation International, Georgetown, Guyana.
- SU, Y.S. & YAJIMA, M. (2015) R2jags using R to run JAGS. *R package version 0.5-7*, 34.
- TROLLE, M. (2003) Mammal survey in the southeastern Pantanal, Brazil. *Biodiversity & Conservation*, 12, 823–836.
- WERNECK, F.P. (2011) The diversification of eastern South American open vegetation biomes: historical biogeography and perspectives. *Quaternary Science Reviews*, 30, 1630–1648.
- WHITNEY, G.G. & DAVIS, W.C. (1986) From primitive woods to cultivated woodlots: Thoreau and the forest history of Concord, Massachusetts. *Journal of Forest History*, 30, 70–81.
- WILKIE, D.S., BENNETT, E.L., PERES, C.A. & CUNNINGHAM, A.A. (2011) The empty forest revisited. *Annals of the New York Academy of Sciences*, 1223, 120–128.
- ZIPKIN, E.F., ROYLE, J.A., DAWSON, D.K. & BATES, S. (2010) Multi-species occurrence models to evaluate the effects of conservation and management actions. *Biological Conservation*, 143, 479–484.